

Product datasheet

anti-Keratin K13 mouse monoclonal, Ks13.1, lyophilized, purified

Short overview

Cat. No.	61007
Quantity	50 µg
Concentration	50 µg/ml after reconstitution with 1 ml dist. water

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	Ks13.1
Immunogen	Keratin K13 of Mr 54,000 purified from human esophagus
Formulation	Lyophilized; reconstitute in 1 ml dist. water (final solution contains 0.09% sodium azide, 0.5% BSA in PBS buffer, pH 7.4)
Synonym	Cytokeratin 13
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage before reconstitution	2-8°C until indicated expiry date
Storage after reconstitution	Up to 3 months at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	ICC/IF, IHC, WB
Reactivity	Bovine, Human, Rat

Applications

Immunocytochemistry (ICC)	Assay dependent
Immunohistochemistry (IHC) - frozen	1:5-1:25 (2 µg/ml - 10 µg/ml)
Immunohistochemistry (IHC) - paraffin	1:5-1:25 (2 µg/ml - 10 µg/ml) (protease treatment and/or microwave treatment recommended)
Western Blot (WB)	1:50-1:100

Background

Ks 13.1 represents an excellent marker to discriminate non-cornified squamous epithelia from those of different origin. Polypeptide reacting: Mr 54,000 polypeptide human keratin K13 (formerly designated cytokeratin 13; with minor affinity to keratins K14, Mr 50,000, and K16, Mr 48,000 and also to K25, formerly designated K25irs1 from inner root sheath of hair follicle). Tumors specifically detected: several squamous cell carcinomas, e.g. cervix carcinoma; transitional cell carcinoma of the bladder; craniopharyngioma. Reactivity on cultured cell lines: cell lines from squamous cell CA, e.g. A-431 from epidermoid CA of vulva; RT 112, RT-4 of urinary bladder CA.


PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com


Product images



Mouse tongue (courtesy of L. Langbein)



IHC analysis of rat tongue using anti-Keratin K13 (Cat. No. 61007). IHC was performed on formalin fixed paraffin embedded sections. The samples were deparaffinized with xylol and ethanol followed by heat induced antigen retrieval with 10 mM citrate buffer. After preparation the tissue was blocked with normal serum for 20 min at RT. The primary antibody anti-Keratin K13 (Cat. No. 61007) was diluted in PBS (1:6, 8 µg/ml) and incubated at 4°C over-night. The secondary antibody biotin anti-mouse was incubated for 30 min at RT. Sections were incubated with ABC solution (VectorLaboratrics) for 30 min at RT. Slides were stained with DAB solution until a brown staining is visible and with Haemalaun for a few minutes. The picture was acquired using microscopy (courtesy of J.Hess, University Hospital Heidelberg).



WB with anti-Keratin K13 antibody (Cat. No. 61007, 1:50), MCF7 whole cell lysate (5-10 ug)

References

Publication	Species	Application
Moll, R. et al. Cytokeratins in normal and malignant transitional epithelium: maintenance of expression of urothelial differentiation features in transitional cell carcinomas and bladder carcinoma cell culture lines. Am. J. Pathol. 132, 123â€“144 (1988).	human	WB,IHC (frozen),IHC (paraffin),ICC-IF
Heid, H. W. et al. Patterns of expression of trichocytic and epithelial cytokeratins in mammalian tissues II. Concomitant and mutually ... Differentiation 37, 215â€“230 (1988).	human	IHC (frozen)
Heid, H. W., Moll, I. & Franke, W. W. Patterns of expression of trichocytic and epithelial cytokeratins in mammalian tissues. I. Human and bovine hair follicles. Differentiation. 37, 137â€“57 (1988).	human	IHC (frozen)