

Product datasheet

anti-AAV VP1/VP2/VP3 mouse monoclonal, B1, liquid, purified, sample

Short overview

Cat. No.	690058S
Quantity	200 µl
Concentration	50 µg/ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	B1
Immunogen	AAV2 capsids
Formulation	PBS, pH 7.4 with 0.09% sodium azide and 0.5% BSA
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	Up to 1 month: 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	Affinity chromatography, Dot blot, ICC/IF, IP, WB
Reactivity	AAV1, AAV2, AAV3, AAV5, AAV6, AAV7, AAV8, AAV9, AAVDJ, AAVrh10

Applications

Affinity Chromatography	Assay dependent
Dot Blot	1:500 (0.1 µg/ml; denaturing conditions)
Immunocytochemistry (ICC)	Assay dependent
Immunoprecipitation (IP)	Assay dependent (precipitation of mainly free VP proteins)
Western Blot (WB)	1:250-1:500 (0.1-0.2 µg/ml)

Background

The B1 antibody reacts with free VP1, VP2 and VP3 of adeno-associated virus (AAV) and at a reduced degree with assembled viral particles. VP1 and VP2 are highly enriched in the nucleus, while non-assembled VP3 is evenly distributed in the nucleus and the cytoplasm. Epitope mapping experiments (Wobus et al., 2000) identified aa726 to aa733 (C-terminus; common to all 3 VP proteins) as the specific binding region. The antibody is also useful for characterization of different stages of infection. Wobus, C. E. et al. Monoclonal antibodies against the adeno-associated virus type 2 (AAV-2) capsid: epitope mapping and identification of capsid domains involved in AAV-2-cell interaction and neutralization of AAV-2 infection. *J. Virol.* 74, 9281-93 (2000).

Limited Use Label License: Research Use Only Product is exclusively licensed to PROGEN Biotechnik GmbH. The use of these products for the PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com

2024 April 26 / Version: 690058S/DS-231121ibg | Page 1


development, manufacturing and sale of secondary products/derivatives which are based on the purchased products and/or which include the purchased product require a royalty based sub-license agreement.

Product images


B1 epitopes in AAV serotypes

AAV1 **K**SANVDF**D**FTVD**N**NGLYTE**P**RP**I**G**T**RYL**T**R**P**L
 AAV2 **K**SVN**V**D**F**FTVD**T**NGVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAV-DJ **K**ST**S**VDF**F**AVN**T**EGVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAV3B **K**SVN**V**D**F**FTVD**T**NGVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAV4 **Q**QNS**L**L**W**AP**D**A**A**G**K**Y**T**E**P**RA**I**G**T**RYL**T**H**L**
 AAV5 **D**P**Q**F**V**D**F**AP**D**ST**G**E**Y**R**T**TR**P**I**G**T**R**Y**L**T**R**P**L**
 AAV6 **K**SANVDF**D**FTVD**N**NGLYTE**P**RP**I**G**T**RYL**T**R**P**L
 AAV7 **K**Q**T**G**V**D**F**AV**D**S**Q**GVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAV8 **K**ST**S**VDF**F**AVN**T**EGVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAV9 **K**SNN**V**E**F**AVN**T**EGVYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAVrh10 **K**ST**N**VDF**F**AVN**T**EGTYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAVhu . 37 **K**ST**N**VDF**F**AVN**T**EGTYSE**P**RP**I**G**T**RYL**T**R**N**L
 AAVrh74 **K**ST**N**VDF**F**AVN**T**EGTYSE**P**RP**I**G**T**RYL**T**R**N**L

Alignment of B1 epitopes in different AAV serotypes.



Western blot analysis of recombinant AAV2 capsid proteins (Cat. No. 640823, 640824, 640825) with B1 antibody (Cat. No. 690058). Western blot analysis was performed on the precise molar ratio of 1:1:10 (VP1:VP2:VP3) either in separate lanes or combined in one lane. The PVDF membranes were blocked with 5% dry milk in PBST (PBS + 0.1% Tween 20) for 1 h at RT. The primary antibody anti-AAV VP1/VP2/VP3 mouse monoclonal, B1 (Cat. No. 690058) was diluted in blocking buffer (antibody concentration 0.5 µg/ml) and incubated for 1 h at RT. The secondary antibody goat anti-mouse IgG HRP was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.



Western blot analysis of recombinant AAV2 capsid proteins (Cat. No. 640823, 640824, 640825) and denatured AAV2 capsids with B1 antibody (Cat. No. 690058). Western blot analysis was performed on 3.3E+09 denatured AAV2 capsids and recombinant AAV2 VP proteins (ratio 1:1:10 - VP1:VP2:VP3). The PVDF membranes were blocked with 5% dry milk in PBST (PBS + 0.1% Tween 20) for 1 h at RT then probed with 0.2 µg/ml (AAV2 capsids) or 0.5 µg/ml (recombinant VP proteins) anti-AAV VP1/VP2/VP3 mouse monoclonal, B1 as primary antibody diluted in blocking buffer for 1 h at 4°C. The secondary antibody goat anti-mouse IgG HRP was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.

References

Publication	Species	Application
Emmanuel, S. N. et al. Structurally Mapping Antigenic Epitopes of Adeno-associated Virus 9: Development of Antibody Escape Variants. J. Virol. 96, (2022).	AAV5, 9	dot blot
Meng, Y. et al. Cell-penetrating peptides enhance the transduction of adeno-associated virus serotype 9 in the central nervous system. Mol Ther Methods Clin Dev. 21, 28-41(2021).	AAV9	IHC/IF
Galibert, L. et al. Functional roles of the membrane-associated AAV protein MAAP. Sci. Rep. 11, (2021).	AAV2	WB
Kuklik, J. et al. Development of a bispecific antibody-based platform for retargeting of capsid modified aav vectors. Int. J. Mol. Sci. 22, 8355 (2021).	AAV2	WB
Zhang, R. et al. Divergent engagements between adeno-associated viruses with their cellular receptor AAVR. Nat. Commun. 10, 3760 (2019).	AAV	WB