

## Product datasheet

anti-Cardiac Actin mouse monoclonal, AC1-20.4.2, liquid, purified, sample

### Short overview

<b>Cat. No.</b>	690075S
<b>Quantity</b>	200 µl
<b>Concentration</b>	50 µg/ml (10 µg)

### Product description

<b>Host</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Isotype</b>	IgG1
<b>Clone</b>	AC1-20.4.2
<b>Immunogen</b>	Synthetic NH2 terminus decapeptide of cardiac isoform of actin
<b>Formulation</b>	PBS pH 7.4 with 0.09% sodium azide and 0.5% BSA
<b>UniprotID</b>	P68034 (Chicken), P68032 (Human), G1STB6 (Rabbit)
<b>Synonym</b>	Actin, alpha cardiac muscle 1, Alpha-cardiac actin [Cleaved into: Actin, alpha cardiac muscle 1, intermediate form], ACTC1, ACTC
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity chromatography
<b>Storage</b>	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
<b>Intended use</b>	Research use only
<b>Application</b>	IHC, WB
<b>Reactivity</b>	Bovine, Chicken, Human, Mouse, Rabbit


### Applications

<b>Immunohistochemistry (IHC) - frozen</b>	1:10 (5 µg/ml; include 0.5 M NaCl in all washing buffers to enhance specificity)
<b>Immunohistochemistry (IHC) - paraffin</b>	1:10 (5 µg/ml; microwave treatment recommended)
<b>Western Blot (WB)</b>	1:1,000 (50 ng/ml; include 1 M NaCl in all washing buffers to enhance specificity)


### Background

Ac1 represents an excellent marker for cardiac tissue; it discriminates fetal (cardiac)  $\alpha$ -actin from all other actin isoforms. Fetal actin can be localized in regenerating skeletal muscle after injury (in satellite cells) and in veins of the umbelical cord. Mab Ac1-20.4.2 shows no cross reaction with other actin isoforms present in skeletal and smooth muscle, provided that stringent experimental conditions have been applied. Polypeptide reacting: Specific for fetal (cardiac) isoform of actin.

## Product images



anti-Cardiac Actin mouse monoclonal, AC1-20.4.2, liquid, purified, sample



Western blot analysis of human heart whole tissue lysate and with anti-Cardiac Actin antibody. Western blot analysis was performed on 20 µg human heart lysate. The PVDF membrane was blocked with 5% milk in PBST (PBS + 0.1% Tween 20) for 1 h at RT. The primary antibody anti-Cardiac Actin mouse monoclonal, AC1-20.4.2 (Cat. No. 690075) was diluted in blocking buffer (antibody concentration 0.05 µg/ml) and incubated for 1 h at RT. The secondary antibody anti-mouse IgG, HRP conjugate was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.

## References

Publication	Species	Application
<a href="#">Domke, L and Franke, W. The cell-cell junctions of mammalian testes..., Cell Tissue Res, 375, 451-482, (2019)</a>	bovine	ICC-IF
<a href="#">Marrocco, V. et al. PKC and PKN in heart disease. J.Mol.Cell.Cardiol. 128, 212-226 (2019)</a>	mouse	WB
<a href="#">Kant, S. et al. Desmoglein 2 mutation provokes skeletal muscle actin expression and accumulation at intercalated discs in murine hearts. J.Cell.Sci. 132, (2019)</a>	mouse	IHC-IF (paraffin)
<a href="#">Hernandez, D. A. et al. Nebulette is a powerful cytolinker organizing desmin and actin in mouse hearts. Mol. Biol. Cell 27, mbc.E16-04-0237 (2016).</a>	mouse	WB
<a href="#">Lindskog, C. et al. The human cardiac and skeletal muscle proteomes defined by transcriptomics and antibody-based profiling. BMC Genomics 16, 475 (2015).</a>	human	IHC