

Product datasheet

AAV6 VP1, recombinant protein

Short overview

Cat. No.	640836
Quantity	10 µg
Concentration	100 µg/ml (1.19 µM)

Product description

Formulation	Liquid, 6 M urea in PBS
Source	Escherichia coli
Molecular Weight	83.6 kDa (calculated Mw from aa sequence)
Purity	> 95% (determined by SDS PAGE)
Product description	N-terminal His-tagged (MGSSHHHHHSSGLVPRGSH) recombinant AAV6 capsid protein VP1
Purification	Ni-NTA chromatography
Storage	-80°C
Intended use	Research use only
Application	Dot blot, SDS PAGE, WB

Applications

Dot Blot	100 ng, depending on primary antibody and detection method
SDS PAGE	1 µg
Western Blot (WB)	5-20 ng, depending on primary antibody and detection method


Background

The AAV capsid consists of three capsid proteins, i.e. VP1, VP2 and VP3, which differ in their N-terminus and encapsulate the genomic ssDNA. In native virus particles, the three proteins form subunits with a ratio of 1:1:10 (VP1:VP2:VP3), in a total number of 60 subunits per capsid. The recombinant AAV6 VP1 protein in combination with recombinant AAV6 VP2 (Cat. No. 640837) and recombinant AAV6 VP3 (Cat. No. 640838) can be used to create a mixture with the precise molar ratio of 1:1:10 to compare the protein composition of the viral capsid in your sample by protein detection methods, e.g. western blot. All three recombinant AAV6 capsid proteins are available as set (Cat. No. 72006) or as individual proteins (Cat. No. 640836, 640837, 640838). Note: please find an example how to prepare western blot samples in the pipetting scheme below. Aliquots of the remaining samples can be stored at -80°C for reuse.


Product images



AAV6 VP1, recombinant protein



Pipetting scheme for western blot analysis using a mix of the AAV6 capsid proteins (Cat. No. 640836, 640837, 640838). To create a VP mixture with the molar ratio 1:1:10 (VP1:VP2:VP3), please pre-dilute VP1 and VP2 1:10 to yield a final concentration of 10 µg/ml (green table). Pipette the pre-diluted VP1 and VP2 proteins and mix them with the undiluted VP3 protein in your sample buffer and water (blue table). The example with 2x and 3x sample buffer and the required volumes are indicated in the pipetting scheme. Thus, in one lane, 10 µl of the VP mix can be loaded onto the SDS PAGE and analyzed by Western blot using the B1 antibody (Cat. No. 690058, Cat. No. 61058-488, Cat. No. 61058-647). Undiluted = 100 µg/ml, pre-diluted = 10 µg/ml



Pipetting scheme for western blot analysis using the AAV6 capsid proteins (Cat. No. 640836, 640837, 640838) in separate lanes. To analyze the molar ratio of 1:1:10, it is recommended to load VP1, VP2 and VP3 as described in the pipetting scheme above. Therefore, the indicated volumes of the proteins (concentration 100 µg/ml) should be diluted with the appropriate amount of sample buffer and distilled water. 10 µl of each solution can be separately loaded onto the SDS PAGE and analyzed by Western blot using the B1 antibody (Cat. No. 690058, Cat. No. 61058-488, Cat. No. 61058-647).

References

Publication	Species	Application
Acevedo, J., et al., Assessment of adeno-associated virus purity by capillary electrophoresis-based western., Mol Ther Methods Clin Dev 32, 101321, (2024).		CE

References

Publication	Species	Application
Acevedo, J., et al., Assessment of adeno-associated virus purity by capillary electrophoresis-based western., Mol Ther Methods Clin Dev 32, 101321, (2024).		CE