

Product datasheet

anti-GST-tag mouse monoclonal, F50-3D12.2, lyophilized, purified, large

Short overview

| | |
|----------------------|---|
| Cat. No. | 910GSTL |
| Quantity | 100 µg |
| Concentration | 0.25 mg/ml after reconstitution with 400 µl PBS |

Product description

| | |
|--------------------------------------|---|
| Host | Mouse |
| Antibody Type | Monoclonal |
| Isotype | IgG1 |
| Clone | F50-3D12.2 |
| Immunogen | Not available or unknown |
| Formulation | Lyophilized; reconstitute in 400 µl sterile PBS, pH 7.4 |
| Conjugate | Unconjugated |
| Purification | Affinity chromatography |
| Storage before reconstitution | 2-8°C until indicated expiry date |
| Storage after reconstitution | -20°C (avoid freeze/thaw cycles) |
| Intended use | Research use only |
| Application | WB |
| Reactivity | GST |


Applications

| | |
|--------------------------|-------------------|
| Western Blot (WB) | 1:500 (0.5 µg/ml) |
|--------------------------|-------------------|

Background

The monoclonal F50-3D12.2 antibody recognizes the GST-tag (Glutathione S-transferase). The GST-tag is commonly added to recombinant proteins and can be used for detection or purification of the tagged protein.

Product images



Western blot analysis of E. coli lysate containing GST-tagged protein with anti-GST-tag antibody. Western blot analysis was performed on 100 ng or 25 ng of E. coli lysate containing GST-tagged protein. Cells were lysed with SDS sample buffer. The PVDF membrane was blocked with 5% dry milk in PBST for 1 h at RT. The primary antibody anti-GST-tag mouse monoclonal, F50-3D12.2 (Cat. No. 910GSTL) was diluted in blocking buffer (antibody concentration 0.5 µg/ml) and incubated for 1 h at RT. The secondary antibody goat anti-mouse IgG polyclonal, HRP conjugate was also diluted in blocking buffer (antibody concentration 0.2 µg/ml) and incubated for 1 h at RT. The bands were visualized by chemiluminescent detection using Pierce™ ECL Western Blotting Substrate.