

## **Product datasheet**

# anti-Laminin rabbit polyclonal, serum

#### Short overview

 Cat. No.
 10765

 Quantity
 250 μl

### **Product description**

Host Rabbit
Antibody Type Polyclonal

Immunogen Laminin isolated from Engelbrecht-Holm-Swarm (EHS)-mouse sarcoma

Formulation Contains 0.09% sodium azide

UniprotID A0A3Q1LUC5 (Bovine),P25391 (Human),P19137 (Mouse),A0A4X1VLN1 (Pig),H0VND7 (Guinea

pig),G1TTH9 (Rabbit),G1TTH9 (Rabbit),D4A409 (Rat)

Synomym Laminin subunit alpha-1, Laminin A chain, Laminin-1 subunit alpha, Laminin-3 subunit alpha,

S-laminin subunit alpha, S-LAM alpha, LAMA1, LAMA

Note Centrifuge prior to opening

ConjugateUnconjugatedPurificationStabilized antiserum

**Storage** Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles

Intended use Research use only

Application IHC, WB

Reactivity Bovine, Human, Mouse, Pig, Rabbit, Rat

### **Applications**

Immunohistochemistry (IHC) - frozen 1:50-1:100

Immunohistochemistry (IHC) - paraffin 1:50-1:100 (proteolytic treatment required; enzyme: pepsin)

Western Blot (WB) Assay dependent

#### Background


Laminin is a glycoprotein (Mr 850 - 1.000 kD, consisting of 3 glycosylated polypeptide chains with molecular weights of 440 and 225 (2x) kD) produced by various human epithelial and mesenchymal cells, and forms an extracellular matrix of thin filaments. In normal tissues, laminin is invariably present in all basal laminas surrounding muscle, nerve, fat and decidua cells and separates epithelial and endothelial cells from abutting connective tissues. Laminin has also been identified within the cytoplasm of breast epithelia, stromal cells of the endometrium, and within endothelial, bile duct epithelial and mesenchymal cells of the liver. Laminin has been found to be involved in cellular activities such as adhesion, spreading, differentiation, polarization, proliferation, locomotion, tissue invasion and chemotactic responses. No cross reaction was obtained with human type I, III, IV and V collagen in immunoblotting, whereas the antibody reacted with a distinct band of appr. 200-220 kD from a 8M Urea extract from amnion basement membrane.

Positive control: skin, kidney.

### **Product images**




anti-Laminin rabbit polyclonal, serum




Chandana, E. P., Maeda, Y., et al. Involvement of the Reck tumor suppressor protein in maternal and embryonic vascular remodeling in mice.

<u>BMC Dev Biol. 2010-08-06.</u> Species/Reactant: Mus musculus (House mouse)Applications: ImmunohistochemistryImage collected and cropped by CiteAb from the following publication, provided under a CC-BY licence.



de Almeida, G. M., Yamamoto, M., et al. Critical roles for murine Reck in the regulation of vascular patterning and stabilization. Sci Rep.

<u>2015-12-11.</u> Species/Reactant: Mus musculus (House mouse)Applications: ImmunohistochemistryImage collected and cropped by CiteAb from the following publication, provided under a CC-BY licence.



de Almeida, G. M., Yamamoto, M., et al. Critical roles for murine Reck in the regulation of vascular patterning and stabilization. Sci Rep.

2015-12-11. Species/Reactant: Mus musculus (House mouse)Applications: Immunohistochemistry-immunofluorescenceImage collected and PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

cropped by CiteAb from the following publication, provided under a CC-BY licence.			
	DOCEN Bistochaile Carbill Mos Cotas Co. 201	D 00400 Heidelberr	

# References

Publication	Species	Application
Domke, L and Franke, W. The cell-cell junctions of mammalian testes, Cell Tissue Res, 375, 451-482, (2019)	bovine	ICC-IF
Costa, B. et al. Intratumoral platelet aggregate formation in a murine preclinical glioma model depends on podoplanin expression on tumor cells. Blood.Adv. 3, 1092-1102 (2019)	rat	IHC (paraffin)
Eisemann, T. et al. Podoplanin expression is a prognostic biomarker but may be dispensable for the malignancy of glioblastoma. Neuro.Oncol. 21, 326-336 (2019)	mouse	IHC (paraffin)
Eisemann, T. et al. An advanced glioma cell invasion assay based on organotypic brain slice cultures. BMC.Cancer. 18, 103 (2018).	mouse	IHC (paraffin)
De Almeida, G. M. et al. Critical roles for murine Reck in the regulation of vascular patterning and stabilization. Sci. Rep. 5, 17860 (2015).	mouse	IHC