

Product datasheet

anti-Keratin K18 mouse monoclonal, Ks18.04, purified, sample (ready-to-use)

Short overview

Cat. No.	61028S
Quantity	600 µl (ready-to-use)

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	Ks18.04
Immunogen	Human keratin K18 from HeLa cytoskeletal preparation
Formulation	PBS pH 7.4 with 0.5% BSA and 0.09% sodium azide
UniprotID	A1XEA5 (Bovine),P05783 (Human),P05784 (Mouse),F1SGG1 (Pig),H0UYZ2 (Guinea pig),Q5BJY9 (Rat),Q5BJY9 (Rat),W5Q5M3 (Sheep)
Synonym	Keratin, type I cytoskeletal 18, Cell proliferation-inducing gene 46 protein, Cytokeratin-18, CK-18, Keratin-18, K18, KRT18, CYK18, PIG46
Conjugate	Unconjugated
Purification	Affinity chromatography
Storage	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	ICC/IF, IHC, WB
Reactivity	Bovine, Dog, Hamster, Human, Mouse, Pig, Rat, Sheep, Trout, Zebrafish

Applications

Immunocytochemistry (ICC)	Assay dependent
Immunohistochemistry (IHC) - frozen	Ready-to-use
Immunohistochemistry (IHC) - paraffin	Ready-to-use (microwave treatment recommended)
Western Blot (WB)	Assay dependent


Background

Ks18.04 represents an excellent marker to discriminate simple epithelia from those of different origin. Tumors specifically detected: all adenocarcinoma; mammary carcinoma, urinary bladder carcinoma, undifferentiated carcinoma, cervix carcinoma, hepatocellular carcinoma. Polypeptide reacting: Mr 45,000 polypeptide (human keratin K18; formerly also designated cytokeratin 18) of all simple type epithelia and basal cells of many squamous, nonepidermal epithelia.


Tested cultured cell lines: MCF-7.



upcyte hepatocytes (courtesy of upcyte technologies GmbH)



IHC of human liver (courtesy of J.Heß, University Hospital Heidelberg)



IHC of rat kidney (courtesy of J.Heß, University Hospital Heidelberg)

References

Publication	Species	Application
Isozaki, Y. et al. The Rho-guanine nucleotide exchange factor Solo decelerates collective cell migration by modulating the Rho-ROCK pathway and keratin networks. Mol Biol Cell. 31, 741-752(2020).	dog	WB
Hojo, M. et al. A histopathological analysis of spontaneous neoplastic and non-neoplastic lesions in aged male RccHan:WIST rats. J.Toxicol.Pathol. 33, 47-55 (2020)	rat	IHC (paraffin)
Santoro, A. et al. p53 Loss in Breast Cancer Leads to Myc Activation, Increased Cell Plasticity, and Expression of a Mitotic Signature with Prognostic Value. Cell.Rep. 26, 624-638.e8 (2019)	mouse	IHC (paraffin)
Norum, J. et al. GLI1-induced mammary gland tumours are transplantable and maintain major molecular features. Int.J.Cancer. , (2019)	mouse	IHC (paraffin)
Ordonez, L. et al. Rapid activation of epithelial-mesenchymal transition drives PARP inhibitor resistance in Brca2-mutant mammary tumours. Oncotarget. 10, 2586-2606 (2019)	mouse	IHC (paraffin)