

Product datasheet

Neurofilament 160 kDa, bovine, 250 µg

Short overview

Cat. No.	62009
Quantity	250 µg

Product description

Source	Bovine spinal cord
Molecular Weight	160 kDa
Isoelectric point	pI 5.1
Purity	> 95% (determined by SDS gelelectrophoresis)
Reconstitution	Reconstitute with 200 µl distilled water (final volume 250 µl). Final solution: 10 mM sodium phosphate, pH 7.5, 2 mM DTT, 6 M urea, 10 mM methylammonium chloride, 1 mM EDTA; Protein concentration: 1 mg/ml (according to Bradford)
Application	Protein standard in 1D and 2D SDS gelelectrophoresis, immunoassays and immunization
Storage	Lyophilized at -2-8°C; reconstituted at -20°C (avoid freeze/thaw cycles)
Intended use	Research use only

Background

Protein standard for immunoblotting, immunization and immunoassays.

Product images



Neurofilament 160 kDa, bovine, 250 µg

References

Publication	Species	Application
Bartos A, Fialova L, Soukupova J, Kukal J, Malbohan I, Pit�ha J: Elevated intrathecal antibodies against the medium neurofilament subunit in multiple sclerosis. J Neurol 254, 20-25 (2007)		
Weber K, Shaw G, Osborn M, Debus E and Geisler N: Neurofilaments, a subclass of intermediate filaments: Structure and expression. Cold Spring Harbor Symp Quant Biol 48, 717 ff (1983)		
Dahl D, Crosby CJ, Gardner EE and Bignami A: Purification of the glial fibrillary acidic protein by anion-exchange chromatography. Analyt Biochem 126, 165 ff (1982)		