

## Product datasheet

### anti-Keratin K19 guinea pig polyclonal, serum

#### Short overview

Cat. No.	GP-CK19
Quantity	100 µl

#### Product description

<b>Host</b>	Guinea pig
<b>Antibody Type</b>	Polyclonal
<b>Immunogen</b>	Human keratin K19, purified from MCF-7 cells
<b>Formulation</b>	Contains 0.09% sodium azide and 0.5% BSA
<b>Stability</b>	Guinea pig (GP68)
<b>UniprotID</b>	P08728 (Bovine), P08727 (Human)
<b>Synonym</b>	Keratin, type I cytoskeletal 19, Cytokeratin-19, CK-19, Keratin-19, K19, KRT19
<b>Note</b>	Centrifuge prior to opening
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Stabilized antiserum
<b>Storage</b>	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
<b>Intended use</b>	Research use only
<b>Application</b>	ICC/IF, IHC, WB
<b>Reactivity</b>	Human
<b>No reactivity</b>	Mouse, Rat

#### Applications

<b>Immunocytochemistry (ICC)</b>	1:100
<b>Immunohistochemistry (IHC) - frozen</b>	1:100
<b>Immunohistochemistry (IHC) - paraffin</b>	1:100 (microwave treatment recommended)
<b>Western Blot (WB)</b>	1:3,000

#### Background

The antiserum represents an excellent marker for all glandular type epithelia, useful for discrimination of glandular epithelial carcinoma from those of different origin. No reaction with hepa-toxic cellular ca.! Polypeptide Reacting: Mr 40,000 polypeptide (keratin K19) of human glandular epithelia.Reactive polypeptide: acidic human keratin K19 (Mr 40,000; formerly also designated cytokeratin 19)Tumors detected: all adeno-carcinoma tested.

#### Product images




© PROGEN in cooperation with Uniklinik Heidelberg, Dr. Jochen Heß

Human head and neck squamous-cell carcinoma (HNSCC) (courtesy of J. Hess, University Hospital Heidelberg)



© PROGEN in cooperation with Uniklinik Heidelberg, Dr. Jochen Heß

Human head and neck squamous-cell carcinoma (HNSCC) (courtesy of J. Hess, University Hospital Heidelberg)



© Lutz Langbein, distributed by PROGEN

Human duodenum (courtesy of L. Langbein)

## References

Publication	Species	Application
<a href="#"><u>Larribere, L. et al. An RNAi Screen Reveals an Essential Role for HIPK4 in Human Skin Epithelial Differentiation from iPSCs. Stem Cell Reports. 9, 1234-1245 (2017).</u></a>	human	IHC-IF (paraffin)
<a href="#"><u>Campard, D., Lysy, P. A., Najimi, M. &amp; Sokal, E. M. Native Umbilical Cord Matrix Stem Cells Express Hepatic Markers and Differentiate Into Hepatocyte-like Cells. Gastroenterology 134, (2008).</u></a>	human	ICC-IF
<a href="#"><u>Langbein, L. et al. Characterization of a Novel Human Type II Epithelial Keratin K1b, Specifically Expressed in Eccrine Sweat Glands. J. Invest. Dermatol. 125, 428â€“444 (2005).</u></a>	human	IHC (frozen)