

Product datasheet

AAV2 empty capsids

Short overview

Cat. No.	66V020
Quantity	100 µl (> 5.0E+11 capsids)
Concentration	> 5.0E+12 capsids/ml; please find the lot-specific concentration on the CoA and on the vial

Product description

Formulation	PBS + 0.014% Tween20 + 1 mM MgCl ₂ + 2.5 mM KCl
Source	Produced in HEK293T cells
Purity	> 95% by SDS-PAGE and silver stain
Quality check	Final titer was assigned on internal reference material using two AAV2 ELISA kits (PRAAV2R and PRAAV2XP); QC included analysis of filling grade and Endotoxin testing
Packaging Plasmid	pDP2rs
Product description	Purified empty adeno-associated virus 2 capsids (AAV2)
Endotoxin	< 1.0 EU/ml
Note	Please centrifuge before opening to ensure complete recovery of vial contents; repeated freeze/thaw cycles and aliquoting can lead to a drop in titer
Purification	Affinity Chromatography (POROS CaptureSelect AAVX Affinity Resin, Thermo Fisher Scientific)
Storage	Up to 2 weeks: 2-8°C; long term storage in aliquots at -80°C; avoid > 5 freeze/thaw cycles
Intended use	Research use only
Application	Dot blot, ELISA, WB

Applications

Dot Blot	Depending on primary antibody and detection method
ELISA	As a positive control in ELISA, dilute in ASSB 1x (provided with PROGENs AAV2 ELISA) and analysis at least in duplicates is recommended
Western Blot (WB)	Depending on primary antibody and detection method

Background

Our AAV2 empty capsids consist of fully assembled, empty AAV2 capsids and have a titer of > 5.0E+12 capsids/ml. Final concentrations are lot-specific and can be found on the corresponding vial. The AAV2 empty capsids are provided with titers above 5.0E+12 capsids/ml in a liquid formulation. Since the buffer does not contain any stabilizing proteins or dyes, the capsids can be used in various applications, including dot blot, western blot and ELISA. The lot-specific titers were assigned using two different AAV2 ELISA (PRAAV2R and PRAAV2XP). Internal standard material calibrated with the ATCC reference standard material for AAV2 (VR-1616) was used for the final titer declaration. Our comprehensive quality control ensures well-characterized capsid material which can be implemented as reference material in a variety of assays to prove the


PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: info@progen.com | Web: www.progen.com


2024 April 25 / Version: 66V020/DS-070823lim | Page 1

validity of the corresponding assay. PROGEN also provides empty capsid material for the AAV serotypes 1, 5, 6, 8 and 9.

Product images



Analytical ultracentrifugation (AUC) of the AAV2 empty capsids shows a strong peak in the 65S range typical for empty capsid with a very small peak of 80-100S corresponding to DNA containing capsids. Capsids were centrifuged at 18000 rpm with 120 scans (1/min) and detected by interference.



SDS PAGE with AAV2 empty capsids. The AAV2 VP1, VP2 and VP3 proteins were separated on a 10% SDS PAGE and visualized by Pierce Silver stain kit (Cat. No. 24612). Only VP1, VP2 and VP3 proteins in the correct stoichiometry of 1:1:10 are detectable indicating a purity of the AAV preparation of > 95%.



AAV2 empty capsids