

## Product datasheet

### Desmin, chicken gizzard, 250 µg

#### Short overview

<b>Cat. No.</b>	62005
<b>Quantity</b>	250 µg

#### Product description

<b>Source</b>	Chicken gizzard
<b>Molecular Weight</b>	53 kDa
<b>Isoelectric point</b>	pI 5.4
<b>Purity</b>	> 98% (determined by SDS gelelectrophoresis)
<b>Reconstitution</b>	Reconstitute with 200 µl distilled water (final volume 250 µl). Final solution: 10 mM sodium phosphate buffer pH 7.5, 6 M urea, 2 mM DTT, 1 mM EDTA, 10 mM methylammonium chloride; Protein concentration: 1 mg/ml (determined by Bradford method).
<b>Application</b>	Protein standard in 1D and 2D SDS gelelectrophoresis, immunoassays and immunization
<b>Storage</b>	Lyophilized at -2-8°C; reconstituted at -20°C (avoid freeze/thaw cycles)
<b>Intended use</b>	Research use only

#### Background

Protein standard for immunoblotting, immunization and immunoassays. Reconstitution to filaments is performed by dissolving in 6 M urea buffer (see above) at concentrations of approx. 0.5 mg/ml. Protofilaments and filament complexes are obtained by dialyzing the resulting polypeptide solution stepwise to a concentration of 4 M urea and then to low salt condition (50 mM NaCl, 2 mM dithiothreitol, 10 mM Tris-HCl, pH 7.4). For immunization purposes, the solution can be further dialyzed against PBS (phosphate buffered saline, e.g. Dulbecco's PBS). - Hatzfeld M and Franke WW (1985). J Cell Biol 101, 1826-1841- Hatzfeld M et al. (1987). J Mol Biol 197, 237-255

#### Product images



Desmin, chicken gizzard, 250 µg