

Product datasheet

anti-Keratin K19 mouse monoclonal, RCK108, supernatant

Short overview

Cat. No.	11417
Quantity	1 ml

Product description

Host	Mouse
Antibody Type	Monoclonal
Isotype	IgG1
Clone	RCK108
Immunogen	Human bladder cell line T24
Formulation	Contains 0.09% sodium azide
UniproID	P08727 (Human)
Synonym	Keratin, type I cytoskeletal 19, Cytokeratin-19, CK-19, Keratin-19, K19, KRT19
Note	Centrifuge prior to opening
Conjugate	Unconjugated
Purification	Hybridoma cell culture supernatant
Storage	Short term at 2-8°C; long term storage in aliquots at -20°C; avoid freeze/thaw cycles
Intended use	Research use only
Application	ICC/IF, IHC, WB
Reactivity	Human, Mammals

Applications

Immunocytochemistry (ICC)	Assay dependent
Immunohistochemistry (IHC) - frozen	1:10-1:20
Immunohistochemistry (IHC) - paraffin	1:10-1:20 (microwave treatment recommended)
Western Blot (WB)	Assay dependent

Background

Mab RCK 108 specifically reacts with keratin K19 (formerly also designated cytokeratin 19; 40 kD protein in immunoblotting), found in most glandular simple type epithelia and tumors derived therefrom (adenocarcinoma), most squamous cell carcinoma. Positive control: Glandular epithelia.

Product images



anti-Keratin K19 mouse monoclonal, RCK108, supernatant

References

Publication	Species	Application
Kwaspen, F. H. et al. Reproducible and highly sensitive detection of the broad spectrum epithelial marker keratin 19 in routine cancer diagnosis. Histopathology 31, 503â€“516 (1997).	human	WB,IHC (paraffin)
Schaafsma, H. E. & Ramaekers, F. C. Cytokeratin subtyping in normal and neoplastic epithelium: basic principles and diagnostic applications. Pathol. Annu. 29 Pt 1, 21â€“62 (1994).	human	WB
Smedts, F. et al. Keratin expression in cervical cancer. Am. J. Pathol. 141, 497â€“511 (1992).	human	IHC (frozen)