

### **Product datasheet**

## anti-MUC5AC/M1 mouse monoclonal, 58M1, purified

#### Short overview

**Cat. No.** 691521

Quantity1 ml (100  $\mu$ g/ml)Concentration100  $\mu$ g/ml

#### **Product description**

HostMouseAntibody TypeMonoclonalIsotypeIgG1 kappaClone58M1

Immunogen Mucin isolated from an ovarian cyst fluid (pure endocervical type according to the Fenoglio's

classification) from an ALeb patient.

Formulation PBS with 0.02% sodium azide
UniprotID P98088 (Human), E9PWB6 (Mouse)

**Synomym** Mucin-5AC, MUC-5AC, Gastric mucin, Major airway glycoprotein, Mucin-5 subtype AC,

Tracheobronchial, Tracheobronchial mucin, TBM, MUC5AC, MUC5

Conjugate Unconjugated

**Purification** Affinity chromatography

Storage 2-8°C

Intended use Research use only Application ELISA, IHC, WB

**Reactivity** Cat, Human, Monkey, Mouse

No reactivity Rat

#### **Applications**

ELISA Assay dependent Immunohistochemistry (IHC) - frozen 1:50-1:100 (1-2 μg/ml)

**Immunohistochemistry (IHC) - paraffin** 1:50-1:100 (1-2 μg/ml; microwave treatment in 10 mM citrate buffer

pH 6.0 recommended)

Western Blot (WB) 1:50-1:100 (1-2 μg/ml)


### Background

58M1 recognizes the peptide core of gastric mucin M1 (now: MUC5AC), and more specifically with the 'e' epitope amongst the a, b, c, d, e, f, g and h protein core epitopes defined by Bara for M1. MUC5AC is present in primary ovarian mucinous cancer and gastric cancer, but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.

PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Positive control: stomach, gastric cancer.

# **Product images**



Gastric cancer