

## Product datasheet

anti-p120 (catenin, delta-1 (TNND1; pTyr96)) mouse monoclonal, EBS-CA-011, purified

### Short overview

<b>Cat. No.</b>	691628
<b>Quantity</b>	1 ml (100 µg/ml)
<b>Concentration</b>	100 µg/ml

### Product description

<b>Host</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Isotype</b>	IgG1 kappa
<b>Clone</b>	EBS-CA-011
<b>Immunogen</b>	Mouse p120 catenin (pY96)
<b>Formulation</b>	PBS with 0.02% sodium azide
<b>UniprotID</b>	O60716 (Human), P30999 (Mouse), D3ZZZ9 (Rat)
<b>Synonym</b>	Catenin delta-1, Cadherin-associated Src substrate, CAS, p120 catenin, p120(ctn, p120(cas, CTNND1, KIAA0384
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity chromatography
<b>Storage</b>	2-8°C
<b>Intended use</b>	Research use only
<b>Application</b>	IHC, WB
<b>Reactivity</b>	Human, Mouse, Rat

### Applications

<b>Immunohistochemistry (IHC) - frozen</b>	1:50-1:100 (1-2 µg/ml)
<b>Western Blot (WB)</b>	1:50-1:100 (1-2 µg/ml)

### Background

The membrane associated protein pp120 Src substrate (p120 catenin, p120cas) was identified as a tyrosine kinase substrate that is phosphorylated in Src transformed cells or in response to growth factor stimulation. It shares structural similarity with the Drosophila Armadillo protein and the vertebrate beta-catenin and gamma-catenin proteins. Its characteristic Arm domain that is composed of 42-amino acid motif repeats evidences this similarity. In the cell, p120 catenin is localized to the E-cadherin/catenins cell adhesion complex. Like beta- and gamma-catenin, p120 catenin directly associates with the cytoplasmic C-terminus of E-cadherin via its Arm domain and may similarly interact with other cadherins. It exists as four isoforms that range in size from 90-115 kDa. Expression of these isoforms is heterogeneous in human carcinomas, suggesting that altered pp120 expression contributes to malignancy due to loss of functional cell adhesions. Multiple tyrosine residues (Y96, Y112, Y228, Y280, Y257, Y291, Y296, and Y302) in p120 catenin are phosphorylated by Src and these phosphorylations may facilitate interaction with PTP1C/SHP-1 in response to EGF stimulation. Thus, p120 catenin is an Arm domain protein that interacts with both cell


PROGEN Biotechnik GmbH | Maaßstraße 30 | D-69123 Heidelberg

Tel.: +49 (0) 6221 8278-0 | Fax: +49 (0) 6221 8278-24 | Email: [info@progen.com](mailto:info@progen.com) | Web: [www.progen.com](http://www.progen.com)

adhesion molecules, such as cadherins and cell signalling molecules, such as PTP1C.

Positive control: carcinoma.

## Product images



Human pancreas