

## Product datasheet

### anti-Cytomegalovirus p65 mouse monoclonal, CMV-227, purified

#### Short overview

<b>Cat. No.</b>	691505
<b>Quantity</b>	1 ml (100 µg/ml)
<b>Concentration</b>	100 µg/ml

#### Product description

<b>Host</b>	Mouse
<b>Antibody Type</b>	Monoclonal
<b>Isotype</b>	IgG1 kappa
<b>Clone</b>	CMV-227
<b>Immunogen</b>	Recombinant CMV-p65 protein
<b>Formulation</b>	PBS with 0.02% sodium azide
<b>Conjugate</b>	Unconjugated
<b>Purification</b>	Affinity chromatography
<b>Storage</b>	2-8°C
<b>Intended use</b>	Research use only
<b>Application</b>	ELISA, FACS, ICC/IF, IHC
<b>Reactivity</b>	CMV

#### Applications

<b>ELISA</b>	Assay dependent
<b>Flow Cytometry (FACS)</b>	0.5-1.0 µg/million cells in 0.1 ml
<b>Immunocytochemistry (ICC)</b>	1:100-1:200 (0.5-1.0 µg/ml)
<b>Immunohistochemistry (IHC) - frozen</b>	1:50-1:100 (1-2 µg/ml)

#### Background

Ten to forty percent of the patients with acquired immunodeficiency syndrome (AIDS) develop cytomegalovirus (CMV) infections. In some patients with AIDS, CMV is detected in the bronchoalveolar lavage fluid (BALF), urine, and other specimens, even when there are no symptoms of CMV disease. An indicator of active CMV infection is needed to facilitate the diagnosis of CMV disease in patients with AIDS or HIV infection. CMV p65 antigen was detected in the leukocytes of both the peripheral blood and BALF during the early phase of CMV disease.

Positive control: CMV-infected cells or tissues.

#### Product images



anti-Cytomegalovirus p65 mouse monoclonal, CMV-227, purified